Acetaminophen (Tylenol) Dosing Chart

| Child's weight (pounds) | 6-11 | 12- 17 | 18- 23 | 24- 35 | 36- 47 | 48- 59 | 60- 71 | 72- 95 | 96+ | lbs |
|---|------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----|------|
| Liquid 160 mg/ 5 milliliters (mL) | 1.25 | 2.5 | 3.75 | 5 | 7.5 | 10 | 12.5 | 15 | 20 | mL |
| Liquid 160 mg/ 1 teaspoon (tsp) | | 1/2 | 3/4 | 1 | 11/2 | 2 | 21/2 | 3 | 4 | tsp |
| Chewable 80 mg tablets | | | 1½ | 2 | 3 | 4 | 5 | 6 | 8 | tabs |
| Chewable 160 mg tablets | | | | 1 | 1½ | 2 | 2½ | 3 | 4 | tabs |
| Adult 325 mg tablets | | | | | | 1 | 1 | 1½ | 2 | tabs |
| Adult 500 mg tablets | | | | | | | | 1 | 1 | tab |

*Source: https://www.stlouischildrens.org/health-resources/dosage-tables/acetaminophen-tylenol-dosage-table

New Tylenol Dissolvable Packets (6 years old and over): 48-71 pounds: 2 packets

72-95 pounds: 3 packets

DO NOT USE UNDER 2 MONTHS OLD.

WHEN TO USE: Treatment of fever or pain.

DOSING: Repeat every 4-6 hours, as needed, but not more than 4 times in 24 hours